

Driven diffusion of vector fields

S. Marculescu¹

Universität-GH Siegen, D-57068 Siegen, Germany

F. Ruiz Ruiz²

*Departamento de Física Teórica I, Facultad de Ciencias Físicas
Universidad Complutense de Madrid, E-28040 Madrid, Spain*

A model for the diffusion of vector fields driven by external forces is proposed. Using the renormalization group and the ϵ -expansion, the dynamical critical properties of the model with gaussian noise for dimensions below the critical dimension are investigated and new transport universality classes are obtained.

PACS numbers: 64.60.Ak, 64.60.Ht, 68.35.Rh.

Keywords: Driven diffusive systems, gaussian noise, renormalization group, universality classes.

¹E-mail: marculescu@physik.uni-siegen.d400.de

²E-mail: ferruiz@eucmos.sim.ucm.es

Fourteen years ago it was suggested that driven diffusive systems may provide suitable models for fast ionic conductors and solid electrolytes [1]. This triggered the interest in such systems, which over the years have become a very good scenario to investigate the properties of far from equilibrium steady states [2]. A simple microscopic realization of a driven diffusive system is an Ising lattice gas in contact with a thermal bath in which an external uniform driving force is introduced [1]. The usual hopping rate to nearest neighbours is modified through a bias in the rates for hopping along the direction of the external field. In a finite lattice with open boundary conditions, the gas evolves to a non-equilibrium steady state with a non-zero particle current. Due to the energy exchange of the system with the heat bath and the driving field, the system also has a steady energy current. So far driven diffusion of scalar quantities, like the excess of transported particles, has been considered. The purpose of this paper is to extend these considerations to vector quantities.

We begin by very briefly recalling the formulation for the standard case of a scalar quantity [3]. Space is taken to be d -dimensional and parameterized by coordinates $(x_i) = (x_{\parallel}, \mathbf{x}_{\perp})$, with $\parallel$ the direction of the driving force and $\perp$ the directions orthogonal to it. The deviation $s(x, t)$ of the density from its uniform average satisfies a continuity equation $\partial_t s + \partial_i J_i(s) = \eta$, with $J_i(s)$ the current generated by ordinary diffusion and the drift in the direction of the driving force, and $\eta(x, t)$ the noise accounting for the heat bath. The contribution of ordinary diffusion to the current is proportional to the gradient of s , while the drift in the $\parallel$ -direction can be estimated, for small density fluctuations s , by expanding in powers of s and retaining terms through order two. This gives the driven diffusion equation [3]

$$\partial_t s - \nu \partial_{\parallel}^2 s - \nu_{\perp} \Delta_{\perp} s + \frac{\lambda}{2} \partial_{\parallel} s^2 = \eta . \quad (1)$$

In the first papers [3] the noise was assumed to be the derivative of a random current, thus conserving the number of particles. In subsequent papers [4] gaussian noise (describing systems with random particle sources and drains and preserving particle number only in the mean) was also considered. For both types of noise it was shown that driven diffusive systems have, for dimensions less than the critical dimension, infrared (IR) stable fixed points at which various quantities scale. Furthermore the corresponding critical exponents have been computed using the ϵ -expansion [3] [6] [4], the values obtained showing that density fluctuations spread faster than diffusively.

It is worth noting that eq. (1) also describes the toppling of grains of sand in a sandpile [6]. As a point of fact, the system described by eq. (1) with gaussian noise was first considered as a model for the hydrodynamic regime of a running sandpile. However one must mention that, after a shift, gaussian noise has vanishing average and this requires particles to be randomly added and removed, whereas in the running sandpile particles

are only added. In other words, gaussian noise does not reproduce the feeding rate of the running sandpile [4]. It is also interesting to mention that for $d = 1$ a field redefinition maps the driven diffusion equation (1) into the Kardar-Parisi-Zhang equation [5], which in turn is identical [2] to the Navier-Stokes equation. This is no longer true for $d \neq 1$, for which the three equations are different.

We now generalize driven diffusive systems to the case where the external forces acting on the system produce drift in a hyperplane of dimension n parameterized by coordinates x_α ($\alpha = 1, \dots, n$). We do this by increasing the number of field variables from 1 to n , so that we have a vector $s_\alpha(x, t)$. The component $s_\alpha(x, t)$ describes the excess of grains transported along the direction x_α . For every component $s_\alpha(x, t)$ we write a continuity equation $\partial_t s_\alpha + \partial_i J_{i\alpha} = \eta_\alpha$. Proceeding now as for the standard case, we obtain

$$\partial_t s_\alpha - \sigma \partial_\beta \partial_\beta s_\alpha - 2\sigma\xi \partial_\alpha \partial_\beta s_\beta - \nu_\perp \Delta_\perp s_\alpha + \frac{\lambda}{2} \partial_\alpha s^2 + \kappa \partial_\beta (n s_\alpha s_\beta - \delta_{\alpha\beta} s^2) = \eta_\alpha, \quad (2)$$

where we have used the notation $s^2 \equiv s_\alpha s_\alpha$. This equation has two anisotropy coefficients, σ and $\sigma\xi$, and two coupling constants, λ and κ . For later convenience, we introduce the coefficient

$$\nu \equiv \sigma(1 + 2\xi). \quad (3)$$

Eq. (2) makes sense for $1 \leq n \leq d$. If $n = 1$, we recover the standard driven diffusion equation (1); if $n = d$, anisotropy is maximal and $\nu_\perp = 0$. We want to investigate the dynamical critical properties of driven diffusive systems based on this equation. In this paper, we concentrate on gaussian noise

$$\begin{aligned} \langle \eta_\alpha(x, t) \rangle &= 0 \\ \langle \eta_\alpha(x, t) \eta_\beta(x', t') \rangle &= 2 \delta_{\alpha\beta} \delta^{(d)}(x - x') \delta(t - t') \end{aligned} \quad (4)$$

and use the renormalization group and the ϵ -expansion to characterize to first order in ϵ the universality classes of the model for dimensions below the critical dimension. A similar program can be carried for conserving noise.

We have already mentioned that eq. (2) may be used to describe drift in a hyperplane. Scenarios in which this may become relevant are oil diffusion and diffusion in disorder media. More generally, eq. (2) may be considered in connection with directed percolation [7]. We recall that in directed percolation, when the probability p of jumping to the nearest neighbour along a preferred direction is set equal to a critical value p_* , a phase transition occurs. At this transition, the system shows scaling behaviour together with non-standard features, like *e.g.* fractal dimensionality for the percolating cluster. One could think of the parameter ξ as playing the role of the probability p , with the difference that ξ is not *a priori* set equal to any critical value but enters the model as one other parameter.

The continuity equation (2) can also be thought of as describing the toppling of grains of sand in an n -dimensional sandpile, with $s_\alpha(x, t)$ the deviation in the direction x_α of the sandpile profile from its uniform average. Since in general $n \neq 1$, one may think of using eq. (2) to describe the hydrodynamic regime of sandpiles with avalanches advancing anisotropically. Yet we must keep in mind that gaussian noise does not describe the feeding rate of a sandpile. Other Langevin equations based on gaussian noise describing surface sandpiles have been proposed in ref. [10]. They differ from (2) in that they double the number of field variables but keep the number of transport directions equal to one. It is important to realize that eq. (2) is not the spherical generalization of the ordinary driven diffusion equation (1) in which, while keeping the number of anisotropy directions equal to one, $s(x, t)$ becomes an n -vector field ($n=2j+1$) carrying an n -dimensional irreducible unitary representation of $O(3)$. For isotropic spherical generalizations of the Navier-Stokes equation and the Kardar-Parisi-Zhan equation, see refs. [8] and [9].

To study the system (2)+(4), we use ref. [11] and recast it as a stochastic quantum field theory model with classical action

$$S_n = \int dt d^d x \pi_\alpha \left[-\pi_\alpha + \partial_t s_\alpha - \sigma \partial_\beta \partial_\beta s_\alpha - 2\sigma \xi \partial_\alpha \partial_\beta s_\beta - \nu_\perp \Delta_\perp s_\alpha + \frac{\lambda}{2} \partial_\alpha s^2 + \kappa \partial_\beta (n s_\alpha s_\beta - \delta_{\alpha\beta} s^2) \right], \quad (5)$$

where $\pi_\alpha(x, t)$ is the response field [12] to the assumed gaussian noise. The action S_n has the following symmetries:

- (a) Translational invariance in all coordinates,
- (b) $O(n) \otimes O(d-n)$ invariance with respect to spatial coordinates,
- (c) Joint reflection of longitudinal coordinates, field variables and response fields,
- (d) Scaling of longitudinal coordinates $(t, x_\alpha, \mathbf{x}_\perp) \rightarrow (t, \varrho x_\alpha, \mathbf{x}_\perp)$, with the anisotropy coefficients, coupling constants and fields transforming as $(\sigma, \xi, \nu_\perp) \rightarrow (\varrho^2 \sigma, \xi, \nu_\perp)$, $(\lambda, \kappa) \rightarrow \varrho^{(n+2)/2} (\lambda, \kappa)$ and $(s_\alpha, \pi_\alpha) \rightarrow \varrho^{-n/2} (s_\alpha, \pi_\alpha)$. Since the parameter ξ is scale invariant, we introduce the invariant coupling constants

$$u \equiv \frac{1}{4\pi} \frac{\lambda}{\sigma^{(n+2)/4}} \quad v \equiv \frac{1}{4\pi} \frac{\kappa}{\sigma^{(n+2)/4}}.$$

In addition, if $n = 1$, the action is invariant under Galilei transformations [4] [6]

$$\begin{aligned} s(x, t) &\rightarrow s(x_\perp, x_\parallel - \lambda at, t) + a \\ \pi(x, t) &\rightarrow \pi(x_\perp, x_\parallel - \lambda at, t). \end{aligned}$$

We note that this symmetry does not generalize to $n > 1$. The reason is that Galilei's invariance requires a constant velocity vector and this is only possible for $n = 1$, since in

this case there is only one preferred direction. In other words, it is precisely the passage from $n = 1$ to $n \neq 1$ what destroys the Galilei symmetry.

The critical dimension of the model is $d_c = 4$. For $d = 4$ the model is ultraviolet renormalizable by power counting, the only primitively divergent one-particle irreducible Green functions being the self-energy $\langle \pi_\alpha(p) s_\beta(-p) \rangle$ and the vertex $\langle \pi_\alpha(p) s_\beta(q) s_\gamma(-p-q) \rangle$, which have linear and logarithmic degrees of divergence respectively. Using dimensional regularization [13], we obtain the following one-loop renormalization group equations:

$$\beta_u \equiv \frac{du}{d \ln \mu} = -u \left(\frac{\epsilon}{2} + \mathcal{H} + n\mathcal{L} + \frac{n+2}{4} \mathcal{F} \right) \quad (6)$$

$$\beta_v \equiv \frac{dv}{d \ln \mu} = -v \left(\frac{\epsilon}{2} + \mathcal{H} + \frac{n+2}{4} \mathcal{F} \right) \quad (7)$$

$$\beta_\xi \equiv \frac{d\xi}{d \ln \mu} = \xi (\mathcal{G} - \mathcal{F}) \quad (8)$$

$$\zeta_\sigma \equiv \frac{d \ln \sigma}{d \ln \mu} = \mathcal{F} \quad (9)$$

$$\zeta_\nu \equiv \frac{d \ln \nu}{d \ln \mu} = \frac{\mathcal{F} + 2\xi\mathcal{G}}{1 + 2\xi}, \quad (10)$$

where μ is the dimensional regularization mass scale, $\epsilon = 4 - d$ is the deviation from the critical dimension and $\mathcal{F}$, $\mathcal{G}$, $\mathcal{H}$ and $\mathcal{L}$ are functions of u , v and ξ given by

$$n(n+2) \mathcal{X}(u, v, \xi) = u^2 x_{uu}(\xi) + uv x_{uv}(\xi) + v^2 x_{vv}(\xi) \quad \mathcal{X} \equiv \mathcal{F}, \mathcal{G}, \mathcal{H}, \mathcal{L}.$$

For completeness we write the expressions of x_{uu} , x_{uv} and x_{vv} :

$$\begin{aligned} f_{uu} &= 0 \\ f_{uv} &= \left[\frac{n^2}{\xi} - n(n+1)(n-2) \right] a_1 - n^2 \left(\frac{1}{\xi} + 2 \right) a_2 - \frac{n}{2} (n+4) a_3 \\ f_{vv} &= \frac{n^2}{2} (-2n^2 + n + 4) - 2n \left(\frac{n}{\xi} + n + 2 \right) a_1 + n^2 \left(\frac{n+2}{\xi} + 4 \right) a_2 - n^2 \left(\frac{n}{\xi} - n + 2 \right) a_3 \\ &\quad - n(n^2 - n - 4) a_4 \\ g_{uu} &= -\frac{n+2}{2\xi} \left[(n-1) a_1 + \left(1 - \frac{n}{4} \right) a_4 \right] \\ g_{uv} &= \frac{n}{2\xi} \left[-\frac{1}{4} (n-1)(n+2)(3n+4) + \left(\frac{n-2}{\xi} - n^2 + 5n + 2 - \frac{8}{n} \right) a_1 \right. \\ &\quad \left. - (n-2) \left(\frac{1}{\xi} + 2 \right) a_2 + (n-1)(n+2) a_3 - \left(\frac{3}{2} n^2 + 4n - \frac{8}{n} \right) a_4 \right] \end{aligned}$$

$$g_{vv} = \frac{n-2}{2n\xi} f_{vv}$$

$$h_{uu} = \frac{2}{\xi} [(n+1)a_1 - na_2 - a_3]$$

$$h_{uv} = \frac{2n}{\xi} \left[n^2 - 2 - (n+1) \left(n - 2 + \frac{4}{n} \right) a_1 - 2(n-2)a_2 + \left(n - 2 + \frac{4}{n} \right) a_3 + 2a_4 \right]$$

$$h_{vv} = -\frac{n}{\xi} \left[4(n^2 - 2) + \left(n^3 - 4n^2 + 2n - \frac{8}{n} \right) a_1 + 2(n-2)^2 a_2 - \left(n^3 + 2n^2 - 6n + 8 - \frac{8}{n} \right) a_3 + 8a_4 \right]$$

$$\ell_{uu} = \frac{1}{\xi} [-(n+3)a_1 + 2a_2 + (n+1)a_3]$$

$$\ell_{uv} = \frac{2n}{\xi} \left[1 + \frac{6}{n} a_1 + \frac{2(n-2)}{n} a_2 + \left(n - 2 - \frac{2}{n} \right) a_3 - (n+1)a_4 \right]$$

$$\ell_{vv} = -\frac{n}{\xi} \left[4 + \left(n - 4 + \frac{12}{n} \right) a_1 - \frac{2(n-2)^2}{n} a_2 + \left(5n - 4 - \frac{4}{n} \right) a_3 - 4(n+1)a_4 \right],$$

with a_1, a_2, a_3 and a_4 taking the form

$$\begin{aligned} a_1 &= a_2 F\left(\frac{n}{2}, 1; 2; \frac{\xi}{1+\xi}\right) & a_2 &= \frac{1}{(1+\xi)^{n/2}} \\ a_3 &= a_4 F\left(\frac{n}{2}, 1; 2; -\frac{\xi}{1+\xi}\right) & a_4 &= \frac{1}{(1+2\xi)^{n/2}} \end{aligned}$$

and $F(a, b; c; z)$ being the hypergeometric function. For $n = 1$ there is no κ -term in S_n , so that $\kappa = 0$ and hence $v = 0$. From this, and using $n = 1$, it follows that $\mathcal{F} = \mathcal{H} + n\mathcal{L} = 0$. Thus $\beta_u = \epsilon/2$ and, as a result, $\beta_u = 0$ at the critical dimension, implying no ultraviolet renormalization of λ . This lack of renormalization holds to all orders in perturbation theory and follows [6] from the non-anomalous character of the Galilei symmetry of the $n = 1$ model mentioned above. For an $n = 1$ driven diffusive system which is classically Galilei invariant but has a quantum mechanical anomaly, see ref. [14].

To characterize the critical behaviour of the model for $d \leq d_c$, we solve the renormalization group equations in the neighbourhood of all IR attractive fixed points (u_*, v_*, ξ_*) . Eqs. (6)-(8) form a system of differential equations describing the dependence of u, v and ξ on μ . Its solution near an IR attractive fixed point gives, upon substitution in eqs. (9)-(10), the μ -dependence of the anisotropy coefficients σ and ν near the fixed point. Indeed, in the neighbourhood of a fixed point the solutions to eqs. (9)-(10) read $\sigma \propto \mu^{\zeta_{\sigma^*}}$ and

$\nu \propto \mu^{\zeta_{\nu*}}$, with

$$\zeta_{\sigma*} \equiv \mathcal{F}|_*, \quad \zeta_{\nu*} \equiv \left. \frac{\mathcal{F} + 2\xi\mathcal{G}}{1 + 2\xi} \right|_*.$$

To have a signal for criticality, at least one of the two exponents $\zeta_{\sigma*}$ and $\zeta_{\nu*}$ must be negative, since σ or ν will then diverge as $\mu \rightarrow 0$. To find the IR attractive fixed point with diverging length scales of the model for $d \leq d_c$, we therefore proceed in two steps:

Step 1. We first find all fixed points by solving the equations $\beta_u = \beta_v = \beta_\xi = 0$ and retain those which are IR attractive. We remind that for a fixed point to be IR attractive, the matrix of derivatives

$$\begin{pmatrix} \beta_{uu} & \beta_{uv} & \beta_{u\xi} \\ \beta_{vu} & \beta_{vv} & \beta_{v\xi} \\ \beta_{\xi u} & \beta_{\xi v} & \beta_{\xi\xi} \end{pmatrix} \quad \beta_{xy} = \left. \frac{\partial \beta_x}{\partial y} \right|_* \quad (11)$$

that results from linearizing eqs. (6)-(8) about the fixed point must have eigenvalues with strictly positive real parts. The case $\xi_* \rightarrow \infty$ must be treated separately, since then there is no renormalization of ξ , hence no equation (8). In this case, fixed points are solutions of $\beta_u = \beta_v = 0$ and the matrix of derivatives of the beta functions is the upper-left 2×2 submatrix of (11).

Step 2. Next we calculate $\zeta_{\sigma*}$ and $\zeta_{\nu*}$ for all the IR attractive fixed points found in step 1 and keep those values which are negative. For ξ_* finite, the condition $\beta_\xi = 0$ and eq. (8) imply that the two critical exponents take the same value:³ $\zeta_{\sigma*} = \zeta_{\nu*}$. For $\xi_* \rightarrow \infty$, however, the condition $\beta_\xi = 0$ is no longer required and $\zeta_{\sigma*}$ and $\zeta_{\nu*}$ may take different values.

Although n has been assumed to take integer values in the range 1 to 4, the results can be analytically continued in n . In fact, analytic continuation gives well-defined beta functions for exotic values of n , like for example $n \rightarrow 0$ or $n \rightarrow -2$. In what follows we take n as a real variable.

In step 1 of this program, one must keep in mind that the values u_* and v_* of the couplings u and v at a fixed point are not physically meaningful. Hence redefining $\bar{u} = \chi u$ and $\bar{v} = \psi v$, with χ and ψ functions of ξ , does not change the physics. Such a redefinition may be used when one or both of u_* and v_* go to infinity to move the fixed point to $\bar{u}_*$ and $\bar{v}_*$ finite. The renormalization group equations for $\bar{u}$ and $\bar{v}$ are straightforward to obtain from eqs.(6)-(8) and involve the functions χ and ψ . The functions χ and ψ must be found such that the IR attractive fixed points of the new renormalization group equations have $\bar{u}_*$ and $\bar{v}_*$ finite. This introduces two new unknowns, χ and ψ , which, together with the complexity of the functions $\mathcal{F}$, $\mathcal{G}$, $\mathcal{H}$ and $\mathcal{L}$, makes it not feasible to analytically carry the

³This also follows from the definitions of ν , ζ_σ and ζ_n given in eqs. (3), (9) and (10).

program above while keeping ξ and n arbitrary. We have thus restricted ourselves to some values of n and ξ that we find of interest. In particular, we have treated the cases

- (i) $n = -2, 0, 1, 2, 3, 4$ with arbitrary (u_*, v_*, ξ_*) , and
- (ii) $\xi_* = 0, (n-2)/2n, \infty$ with arbitrary (n, u_*, v_*) ,

allowing in both instances for redefinitions of u and v as described above. The values $\xi_* = 0$ and $\xi_* = \infty$ considered in case (ii) correspond respectively to $\pi_\alpha \partial^2 s_\alpha$ and $\pi_\alpha \partial_\alpha \partial_\beta s_\beta$ being the dominant kinetic anisotropic term in the classical action S_n . The idea of looking for critical points at $\xi_* = (n-2)/2n$ stems from the simple relation between f_{vv} and g_{vv} for arbitrary n , namely $2n\xi g_{vv} = (n-2)f_{vv}$. Because of this relation, the equation $\beta_\xi = 0$ simplifies considerably if $\xi_* = (n-2)/2n$, thus making possible to find IR attractive fixed points with finite ξ_* . Another motivation to consider $\xi_* = (n-2)/2n$ is to investigate whether the passage from $n \neq 1$ to $n = 1$ is smooth, since the case $n = 1$ can be recovered from $\xi_* = -1/2$ and $\sigma \rightarrow \infty$.

The results that we have obtained for (i) and (ii) are as follows (we provide the values of the physical quantities n , $\zeta_{\nu*}$ and $\zeta_{\sigma*}$, which characterize the universality classes). For

n	ξ_*	$\zeta_{\nu*}$	$\zeta_{\sigma*}$
3	1/6	-0.1257ϵ	-0.1257ϵ
	∞	positive	-0.2848ϵ
	∞	-0.1068ϵ	-0.4726ϵ
4	1/4	$-158\epsilon/629$	$-158\epsilon/629$
	∞	positive	-0.1324ϵ
	∞	-0.3731ϵ	-0.3526ϵ

Table 1: Universality classes for $n = 3, 4$.

case (i) there are no IR attractive fixed points with diverging length scales if $n = -2, 0, 2$. If $n = 1$, the results of ref. [6] are recovered. And if $n = 3, 4$, we obtain the universality classes in Table 1. For case (ii) we obtain the universality classes:

- $\xi_* = (n-2)/2n$, with $n > 2$ and critical exponents

$$\zeta_{\sigma*} = \zeta_{\nu*} = - \frac{2f_{vv}\epsilon}{(n+2)f_{vv} + 4h_{vv}} \Big|_{\xi=(n-2)/2n}. \quad (12)$$

Note that the critical exponent is a complicated but well-defined function of n .

- $\xi_* = \infty$, with $n = n_0 \equiv (1 + \sqrt{33})/4$ and critical exponents

$$\zeta_{\nu*} = 0 \quad \zeta_{\sigma*} = -\epsilon. \quad (13)$$

- $\xi_* = \infty$, with $n > 2$ and critical exponents

$$\zeta_{\sigma*}^{\pm} = \frac{1}{2} n b_1 \epsilon \frac{2(n+1) X_{\pm} + n b_3}{X_{\pm} (b_1 b_5 - b_2 b_4) + b_1 b_6 - b_3 b_4} \quad (14)$$

$$\zeta_{\nu*}^{\pm} = \frac{1}{16} (n-1) \epsilon \frac{X_{\pm} [n(3n+4)(n-2)b_1 - 8b_2] - 8b_3}{X_{\pm} (b_1 b_5 - b_2 b_4) + b_1 b_6 - b_3 b_4}, \quad (15)$$

where

$$\begin{aligned} b_1 &= -\frac{1}{n-2} (3n^2 + 7n - 2) & b_4 &= -\frac{1}{n-2} (n^2 - 3n - 6) \\ b_2 &= -\frac{1}{8} n (3n+1) (n^2 + 2n - 12) & b_5 &= \frac{1}{2} n (3n^2 - 3n - 10) \\ b_3 &= \frac{1}{2} (2n^2 - n - 4) & b_6 &= -\frac{1}{8} n^2 (n+2) (2n^2 - n - 4) \end{aligned}$$

and

$$X_{\pm} = \frac{1}{2b_1} (-b_2 \pm \sqrt{b_2^2 - 4b_1 b_3}).$$

The exponents $\zeta_{\nu*}^-$ and $\zeta_{\sigma*}^{\pm}$ are negative for all $2 < n \leq 4$, whereas $\zeta_{\nu*}^+$ is negative only for $2 < n < 2.0060$. It is worth noting that the universality classes for $n = -2, 0, 2, 3, 4$ are particular cases of eqs. (12)-(15).

The anomalous dimensions $\zeta_{\nu*}$ and $\zeta_{\sigma*}$ found above are related to power laws in physical correlation functions and therefore should be directly measurable. In accordance with the continuity equation (2) and owing to the vector character of s_{α} , physical correlators are expectation values of products of tensor currents

$$J_{\alpha\beta} = -\sigma \partial_{\alpha} s_{\beta} - 2\sigma \xi \partial_{\beta} s_{\alpha} + \frac{\lambda}{2} \delta_{\alpha\beta} s^2 + \kappa (n s_{\alpha} s_{\beta} - \delta_{\alpha\beta} s^2).$$

Since $J_{\alpha\beta}$ contains linear and quadratic terms in s_{α} , the computation of physical correlators involves renormalization of composite operators. For $n = 1$, due to the non-renormalization of the coupling constant, renormalization of composite operators can be fully eliminated by considering correlators quadratic in the current [4]. For $n \neq 1$, since the coupling constants do renormalize, correlators cubic in $J_{\alpha\beta}$ must also be considered. The analysis of physical correlators and their relation to the critical exponents $\zeta_{\nu*}$ and $\zeta_{\sigma*}$ is in progress and will be presented elsewhere.

We conclude with a few comments:

Comment 1. The anisotropy dimension n at which the critical exponents $\zeta_{\nu*}$ and $\zeta_{\sigma*}$ occur is in general non-integer. It is particularly noticeable that for $\bar{u}_*^2 = 0$, $\bar{v}_*^2$ finite and $\xi_* = \infty$, the only IR attractive fixed point with diverging length scales is that given in

eq. (13) and occurs at $n = (1 + \sqrt{33})/4$. With the caution due to the one-loop nature of our results, this suggests a fractal behaviour of the model. The similarity in some regards of our model with directed percolation may provide some insight in understanding these points.

Comment 2. The fixed points associated with the critical exponents in eqs. (12)-(13) have either $\bar{u}_* = 0$ or $\bar{v}_* = 0$, while the fixed points associated with the critical exponents in eq. (14)-(15) have $\bar{u}_*, \bar{v}_* \neq 0$. We have not been able to find IR attractive fixed points with all three parameters $\bar{u}_*$, $\bar{v}_*$ and ξ_* finite and non-zero. Our results show that if such a point exists to first order in ϵ , it has fractal dimension.

Comment 3. Eqs. (14)-(15) give for $n > 2$ two sets of critical exponents at $\xi_* = \infty$ which are associated with two different attractive IR fixed points. This is so since $\bar{u}_*^2$ and $\bar{v}_*^2$ are two-valued functions of ξ for $\xi \rightarrow \infty$ and $\bar{u}_*, \bar{v}_* \neq 0$. Depending on the configuration of all fixed points, the system will flow in parameter space to one or another IR fixed point and will reach one or another set of critical exponents.

Comment 4. If we take $\lambda = 0$ in S_n , we find universality classes at $\xi_* = \infty$ for $n > n_0$, with critical exponents $\zeta_{\nu*} = 0$ and $\zeta_{\sigma*} = -2\epsilon/(n + 2)$. On the other hand, if we set $\kappa = 0$, we obtain universality classes at $\xi_* = \infty$ for $n > 2$, with critical exponents $\zeta_{\nu*} = 0$ and $\zeta_{\sigma*} = -\epsilon/6$. Note however that, despite the simplicity and beauty of these results, for $\lambda = 0$ or $\kappa = 0$ perturbation theory may generate at higher loops divergent relevant operators which are not in the classical action S_n . For this reason, we have considered the more general case of two coupling constants.

Acknowledgment. The authors are grateful to Bo Zheng for many clarifying discussions. FRR is grateful to the Institute for Theoretical Physics in Heidelberg for its hospitality and to the Alexander von Humboldt Foundation for support through a Research Fellowship.

References

- [1] S. Katz, J.L. Lebowitz and H. Spohn, Phys. Rev. **B28** (1983) 1655; J. Stat. Phys. **34** (1984) 497.
- [2] B. Schmittmann and R.K.P. Zia, *Statistical mechanics of driven diffusive systems*, vol. 17 of the collection *Phase transitions and critical phenomena*, edited by C. Domb and J.L. Lebowitz (Academic Press, London 1995).
- [3] H.K. Janssen and B. Schmittmann, Z. Phys. **B63** (1986) 517; **B64** (1986) 503.

- [4] V. Becker and H.K. Janssen, Phys. Rev. **E50** 1114 (1994).
- [5] M. Kardar, G. Parisi and Y.C. Zhang, Phys. Rev. Lett. **56** (1986) 889.
- [6] T. Hwa and M. Kardar, Phys. Rev. Lett. **62** 1813 (1989); Phys. Rev. **A45** 7002 (1992).
- [7] D. Stauffer and A. Aharony, *Percolation theory*, (Taylor & Francis, London 1994).
- [8] C.Y. You and P. Weichman, Phys. Rev. Lett. **70** (1993) 1101.
- [9] J.P. Doherty, Phys. Rev. Lett. **72** (1994) 2041.
- [10] J.P. Bouchard, M.E. Cates, J. Ravi Prakash and S.F. Edwards, J. Phys. I France **4** 1383 (1994).
A. Mehta, J.M. Luck and R.J. Needs, Phys. Rev. **E53** 92 (1996).
P. Biswas, A. Majumdar, A. Mehta and J.K. Bhattacharjee, *Some properties of a class of coupled Langevin equations*, cond-mat/9708155.
- [11] H.K. Janssen, Z. Phys. **B23** (1976) 377.
C. De Dominicis, J. Phys. (Paris) Colloq. **37** (1976) C-247.
- [12] P.C. Martin, E.D. Siggia and H.H. Rose, Phys. Rev. **A8** (1973) 423.
- [13] G. 't Hooft and M. Veltman, Nucl. Phys. **B44** 189 (1972).
- [14] H.K. Janssen, Phys. Rev. Lett. **78** 1082 (1997).